















## 5.0 REFERENCES

- [1] J.W. Tester, M. Modell, *Thermodynamics and its Applications*, 3rd ed., Prentice Hall PTR, Upper Saddle River, N.J., 1997.
- [2] F. García-Sánchez, J. Schwartzenruber, M.N. Ammar, H. Renon, Modeling of multiphase liquid equilibria for multicomponent mixtures, *Fluid Phase Equilibria*, 121 (1996) 207-225.
- [3] J.M. Prausnitz, A. Firoozabadi, J. Wu, *Molecular-Thermodynamic Framework for Asphaltene-Oil Equilibria*, *American Institute of Chemical Engineers* 44, (1998) 11.
- [4] J.W. Cahn, J.E. Hilliard, Free energy of a nonuniform system. III. Nucleation in a two-component incompressible fluid, *J. Chemical Physics.*, 31 (1959) 688-699.
- [5] J.W. Cahn, J.E. Hilliard, Spinodal decomposition: reprise, *Acta Metallurgica.*, 19 (1971) 151-161.
- [6] T. Hashimoto, Dynamics in spinodal decomposition of polymer mixtures, *Phase Transitions*, 12 (1988) 47-119.
- [7] E. Kiran, Polymer miscibility and kinetics of pressure-induced phase separation in near-critical and supercritical fluids, *NATO Science Series I E*, 366 (2000) 167-192.
- [8] R. Bubbico, E. Salzano, Acoustic analysis of blast waves produced by rapid phase transition of LNG released on water, *Safety Science*, 47 (2009) 515-521.
- [9] N.A. Darwish, S.A. Al-Muhtaseb, A comparison between four cubic equations of state in predicting the inversion curve and spinodal curve loci of methane, *Thermochimica Acta*, 287 (1996) 43-52.
- [10] J.M. Smith, H.C. Van Ness, M.M. Abbott, *Introduction to chemical engineering thermodynamics*, McGraw-Hill, 2005.
- [11] J.P. Wauquier, *Petroleum Refining-Separation Processes*, Editions TECHNIP, Paris, 2000
- [12] M. Ko, S. Na, S. Kwon, S. Lee, H. Kim, Liquid-Liquid Equilibria for the Binary Systems of N-Formylmorpholine with Branched Cycloalkanes, *J. Chemical and Engineering Data*, 48 (2003) 699-702.
- [13] H. Matsuda, A. Kitabatake, M. Kosuge, K. Kurihara, K. Tochigi, K. Ochi, Liquid-liquid equilibrium data for binary perfluoroalkane (C<sub>6</sub> and C<sub>8</sub>)+n-alkane systems, *Fluid Phase Equilibria*, 297 (2010) 187-191.
- [14] K. Kurihara, Y. Yamanaka, H. Matsuda, K. Tochigi, K. Ochi, T. Furuya, Determination and correlation of liquid-liquid equilibria for nine binary acetonitrile+n-alkane systems, *Fluid Phase Equilibria*, 302 (2011) 109-114.
- [15] J. Yang, C. Peng, H. Liu, Y. Hu, Liquid-liquid equilibria of polymer solutions with oriented interactions, *Fluid Phase Equilib.*, 249 (2006) 192-197.
- [16] B.H. Chang, Y.C. Bae, Liquid-liquid equilibria of binary polymer solutions with specific interactions, *Polymer*, 39 (1998) 6449-6454.
- [17] C. Browarzik, D. Browarzik, Liquid-liquid equilibrium calculation in binary water+nonionic surfactant C<sub>i</sub>E<sub>j</sub> systems with a new mass-action law model based on continuous thermodynamics, *Fluid Phase Equilibria*, 235 (2005) 127-138.